



J. Plankton Res. (2016) 38(3): 393–400. First published online April 6, 2016 doi:10.1093/plankt/fbw023

HORIZONS

Metabarcoding of marine zooplankton: prospects, progress and pitfalls

ANN BUCKLIN^{1*}, PENELOPE K. LINDEQUE², NAIARA RODRIGUEZ-EZPELETA³, AITOR ALBAINA⁴ AND MAIJU LEHTINIEMI⁵

¹DEPARTMENT OF MARINE SCIENCES, UNIVERSITY OF CONNECTICUT, GROTON, CT 06340, USA, ²PLYMOUTH MARINE LABORATORY, PLYMOUTH PL1 3DH, UK, ³MARINE RESEARCH DIVISION, AZTI, SUKARRIETA, BIZKAIA 48395, SPAIN, ⁴DEPARTMENT OF GENETICS, PHYSICAL ANTHROPOLOGY AND ANIMAL PHYSIOLOGY, UNIVERSITY OF THE BASQUE COUNTRY UPV/EHU, LEIOA 48940, SPAIN AND ⁵MARINE RESEARCH CENTER, FINNISH ENVIRONMENT INSTITUTE, 00251 HELSINKI, FINLAND

*CORRESPONDING AUTHOR: ann.bucklin@uconn.edu

Received October 10, 2015; accepted February 28, 2016

Corresponding editor: Roger Harris

Metabarcoding (large-scale taxonomic identification of complex samples via analysis of one or few orthologous DNA regions, called barcodes) is revolutionizing analysis of biodiversity of marine zooplankton assemblages. Metabarcoding relies on high-throughput DNA sequencing (HTS) technologies, which yield millions of DNA sequences in parallel and allow large-scale analysis of environmental samples. Metabarcoding studies of marine zooplankton have used various regions of nuclear small- (18S) and large-subunit (28S) rRNA, which allow accurate classification of novel sequences and reliable amplification with consensus primers, but—due to their relatively conserved nature—may underestimate species diversity in a community. To discriminate species, more variable genes are needed. A limited number of metabarcoding studies have used mitochondrial cytochrome oxidase I (COI), which ensures detection of species-level diversity, but may require group-specific primers and thus result in inconsistent amplification success rates. Reference databases with sequences for accurately-identified species are critically needed to allow taxonomic designation of molecular operational taxonomic units (MOTU) and comparison with previous studies of zooplankton diversity. Potential and promising applications of metabarcoding include rapid detection of impacts of climate change, monitoring and assessment of ecosystem health, calculation of biotic indices, characterization of food webs and detection of introduced, non-indigenous species.

KEYWORDS: metabarcoding; DNA barcoding; biodiversity; high-throughput DNA sequencing; zooplankton

INTRODUCTION

Marine zooplankton are rapid-responders to environmental variation associated with regime shifts and climate change, which may cause significant and potentially

accelerating losses in species diversity (Beaugrand *et al.*, 2010; Möllmann and Diekmann, 2012). However, the systematic complexity of the zooplankton assemblage, with numerous cryptic and sibling species, and the lack of

diagnostic characters for immature (larval) stages constitute important impediments to our understanding of global-to-local patterns of biodiversity and biogeography. In recent years, high-throughput DNA sequencing (HTS) technologies have resulted in dramatic advances in practical, cost-effective molecular approaches to the analysis of environmental samples (Bourlat *et al.*, 2013; Ji *et al.*, 2013). In particular, metabarcoding (i.e. the large-scale taxonomic identification of a complex sample via analysis of one or few orthologous DNA regions, called barcodes) has the significant advantage of detecting the ‘hidden diversity’ of zooplankton assemblages, including mero-, holo- and ichthyoplankton (Lindeque *et al.*, 2013). Metabarcoding of the pelagic assemblage is yielding new insights into marine biodiversity, as most marine species (including fish, macro-invertebrates etc.) are planktonic at some point in their life cycle.

The metabarcoding approach involves a variety of laboratory and data analysis steps upon which biodiversity estimates rely (Fig. 1). First, the DNA present in the sample (which can be intracellular or extracellular) is extracted from the whole sample; second, the barcode of choice is amplified using consensus or taxonomic group-specific PCR primers and appropriate amplification conditions; third, the PCR products are sequenced on an HTS platform (each nucleotide sequence is called a ‘read’); and finally, the obtained sequences are processed for quality control, grouped in molecular operational taxonomic units (MOTU), and compared with a reference database for taxonomic assignment. The metabarcoding approach, including these steps and their implications for biodiversity assessments of multicellular zooplankton, is the subject of this article.

CHOICE OF BARCODE

Metabarcoding studies of zooplankton assemblages have used a number of marker gene regions to characterize biodiversity patterns across different systematic levels and to address specific hypotheses. To date, the most frequently used gene regions are portions of the nuclear small-subunit ribosomal RNA gene (18S rRNA), which shows consistent patterns of divergence across invertebrate and vertebrate taxa, and discriminates genera, families and higher taxonomic groups (Mallatt *et al.*, 2004). The V9 hypervariable region of 18S rRNA was developed as a standard marker of marine microbial eukaryotic diversity (Amaral-Zettler *et al.*, 2009) and has also been used for analysis of zooplankton assemblages (Pearman *et al.*, 2014; De Vargas *et al.*, 2015; Pearman and Irigoien, 2015; Albaina *et al.*, 2016). Several additional hypervariable regions of the 18S rRNA gene have been used for

zooplankton metabarcoding studies, including V1–V2 (Lindeque *et al.*, 2013); V4 (Sun *et al.*, 2015) and V7–V9 (Hirai *et al.*, 2015b). These studies allow examination of impacts of both different gene regions and also different sequence lengths, with the latter determined largely by the constraints of the HTS platform and associated protocols. Zooplankton metabarcoding studies have also used portions of the nuclear large-subunit 28S rRNA (Hirai *et al.*, 2013, 2015a; Hirai and Tsuda, 2015). These nuclear rDNA regions have allowed accurate classification of novel sequences and reliable amplification with consensus primers, but due to the relatively conserved nature of this gene may underestimate the diversity of species in a community (Tang *et al.*, 2012).

Analysis of species-level diversity and distribution of marine zooplankton using metabarcoding approaches will require development of reliable HTS protocols for more variable genes. The mitochondrial cytochrome oxidase I (COI) barcode region (Hebert *et al.*, 2003) is one of the most commonly sequenced regions for analysis of species diversity among marine animals (Bucklin *et al.*, 2011), including zooplankton (Bucklin *et al.*, 2010a, b). A limited number of metabarcoding studies of zooplankton species biodiversity based on COI sequences has been published (Machida *et al.*, 2009; Bourlat *et al.*, 2013; Zaiko *et al.*, 2015b). Another mitochondrial gene frequently used for identification and discrimination of zooplankton species is mitochondrial 16S rRNA (Lindeque *et al.*, 1999, 2006; Goetze, 2010), which is considered by some to be a more reliable marker, especially for cnidarians (Zheng *et al.*, 2014; Lindsay *et al.*, 2015). These studies confirm the usefulness of both mitochondrial genes for accurate species identification and discrimination, but also clearly demonstrate that metabarcoding approaches for detection of species-level diversity face significant technical challenges, including the need for cocktails of group-specific primers (Bucklin *et al.*, 2010b), and consequent inconsistent amplification success rates among the various taxonomic groups of marine zooplankton.

REFERENCE BARCODE DATABASE FOR SPECIES IDENTIFICATION

Metabarcoding-based species identification requires taxonomically complete and geographically comprehensive reference databases of DNA sequences for each species and for all gene regions. Absolutely essential is that reference specimens are accurately identified to species by a recognized expert taxonomist; inaccurate and incomplete identifications remain a persistent impediment to the use of metabarcoding for analysis of species-level zooplankton




Fig. 1. Schematic representation of the taxonomic analysis of a zooplankton sample using morphological identification (left arrows), barcoding (middle arrows) or metabarcoding (right arrows). Figure modified from Corell and Rodriguez-Ezpeleta (Corell and Rodriguez-Ezpeleta, 2014).

biodiversity. A reference database can also serve as a valuable resource for researchers to confirm species identifications when morphological taxonomic expertise is limited and when training new taxonomists.

Reference databases are growing for several of the gene regions most frequently used for zooplankton metabarcoding. Most notable are the comprehensive SILVA databases

(<http://www.arb-silva.de/>) with aligned small- (18S) and large-subunit (28S) rRNA sequences for all three domains of life (Quast *et al.*, 2013). Also noteworthy is the archive of COI sequences for thousands of species of marine animals now available in public repositories, including the GenBank Barcode of Life section (<http://www.ncbi.nlm.nih.gov/genbank/barcode>) and the Barcode of Life Database

(<http://www.boldsystems.org>), which provide a valuable reference library that has been likened to a Rosetta Stone for species identification (Bucklin *et al.*, 2010b).

COMPARISONS OF METABARCODING AND MORPHOLOGICAL MEASURES OF ZOOPLANKTON BIODIVERSITY

With the potential for newly emerging metabarcoding analyses to overtake, and perhaps be used as an alternative to morphological analysis in measuring zooplankton diversity, it is critical to compare and contrast the two approaches. While metabarcoding can provide a broad assessment of zooplankton diversity and taxon richness, we should not oversell this relatively new technique in its current state, nor be hasty in replacing morphological analysis, since both techniques have their costs and benefits.

Morphological analysis of zooplankton results in numerical abundance, with a possibility to convert to biomass. HTS technologies provide the ability to read millions of DNA sequences in parallel, making them ideally suited for large-scale biodiversity analyses of samples (Shokralla *et al.*, 2012). To make sense of the data, the sequences are usually clustered into MOTUs (Floyd *et al.*, 2002), based on a similarity threshold (Fonseca *et al.*, 2010). When defining the similarity threshold, both barcode length and intra- and inter-specific degree of conservation need to be taken into account (Lindeque *et al.*, 2013; Brown *et al.*, 2015; Hirai *et al.*, 2015a). Care must be taken in construction of the MOTUs to obtain the most relevant and realistic assessment of diversity and species richness. If the similarity threshold used to cluster sequences into MOTUs is too high, this may lead to an overestimation of taxon richness; however, if the threshold is too low, it is likely that the taxon richness will not align to species richness. Thus, a critical challenge in using metabarcoding to estimate biodiversity is to examine the relationship between MOTU number and species richness (Carugati *et al.*, 2015).

Taxonomic identification of MOTUs can best be made by comparing a representative sequence from each MOTU against a nucleotide database. However, annotation must only be made against well-populated reference libraries based on correctly-identified specimens (see above). In summary, metabarcoding provides a number of sequence reads that can be clustered into MOTUs and with care and consideration can provide estimations of taxon richness that may approximate species richness.

While some pitfalls of metabarcoding, such as lack of identification of individual life stages (since DNA sequences will be identical for eggs, larvae, adults or other developmental stages of a given species), will be nearly impossible to overcome, metabarcoding does present promises above those of traditional morphological methods.

Metabarcoding can discriminate spatial and temporal patterns of variability in planktonic assemblages (Eiler *et al.*, 2013; Massana *et al.*, 2015). Analysis of zooplankton assemblages with metabarcoding has revealed previously hidden taxonomic richness, especially for hard-to-identify meroplankton (e.g. bivalves, gastropods, polychaetes), rare species and parasites in comparison with morphological analysis (Lindeque *et al.*, 2013). Recent metabarcoding studies, such as the TARA oceans expedition (de Vargas *et al.*, 2015) with huge spatial-scale sampling, great depth of sequencing and comprehensive taxonomic analysis, revealed that MOTU diversity is likely to be much higher than described species of marine eukaryotic plankton, especially in the smaller organismal size fraction. As metabarcoding progresses, continued support for traditional morphological analysis will remain critically important to allow direct comparison between morphological and molecular approaches and to gain better understanding of how barcode choice, length and intra- and inter-specific variation influence the similarity threshold. Such studies will ensure that metagenetic assessment of diversity and species richness is both realistic and relevant.

CHALLENGES OF QUANTIFICATION OF TAXON ABUNDANCE OR BIOMASS USING METABARCODING

The sensitivity of metabarcoding analysis to detect and discriminate rare and cryptic species has been widely reported, including for zooplankton communities (Zhan and MacIsaac, 2015). Although presence/absence is critical for biodiversity monitoring, quantification of taxon abundance or relative abundance is needed for community characterization and for the assessment of many biological indices (Bourlat *et al.*, 2013; Aylagas *et al.*, 2014). Quantification of relative abundances of taxa above the species level has been shown to match morphological analyses for some zooplankton groups and samples (Lindeque *et al.*, 2013; Hirai *et al.*, 2015b), but metabarcoding analysis has not shown good agreement with species abundance data from morphological taxonomic analysis (Mohrbeck *et al.*, 2015). In one study (Sun *et al.*, 2015), a general trend was detected that low-abundance species usually corresponded to low-abundance sequence reads; however, the authors urged caution when using HTS-based approaches to make quantitative inferences.

The number of sequencing reads associated with a MOTU can approximate to biomass (Lindeque *et al.*, 2013) and has been shown to correlate with dry weight of the taxon (Hirai *et al.*, 2015b). It is likely that the correlation between biomass and sequencing reads is not linear and is affected by various biases introduced at different stages, e.g. DNA extraction, PCR amplification, DNA pooling and

bioinformatics sorting (Bik *et al.*, 2012). Among possible approaches to address such biases are comparative analysis of RNA (which ensures detection only of living organisms) and approaches that do not require initial PCR amplification (Dowle *et al.*, 2015). An important challenge for quantification using metabarcoding is that multicopy genes, such as ribosomal and mitochondrial genes, vary in copy number across different animal taxa (Prokopowich *et al.*, 2003). It may eventually be possible to calibrate bias due to gene copy number variation (CNV) using low-diversity and/or mock samples, perhaps using quantitative PCR (qPCR) to determine gene copy numbers (Amend *et al.*, 2010). Until these issues are resolved, metabarcoding will remain a semi-quantitative method for biodiversity analysis.

APPLICATIONS OF METABARCODING FOR ECOSYSTEM MONITORING AND MANAGEMENT

Potential applications of metabarcoding in marine monitoring include calculation of biotic indices based on taxonomic composition, characterization of trophic interactions and food web structure and detection of non-indigenous species (NIS). Although applications of metabarcoding for environmental monitoring seem quite straight-forward and work in theory, the routine implementation of this approach still requires the development of standardized practices at each step of the procedure (Aylagas *et al.*, 2014). Because some marine monitoring-related indices, e.g. AZTI's Marine Biotic Index (Aylagas *et al.*, 2014), rely on the presence of taxa that are either sensitive to or tolerant of pollution, the ability to detect all organisms present in the sample is crucial. Careful evaluation of the accuracy of the taxonomic composition inferred from metabarcoding is necessary before this method can be regularly applied to assess ecosystem status.

Another promising frontier for metabarcoding is the analysis of environmental DNA (eDNA) or free DNA molecules that are present outside of organisms (Bohmann *et al.*, 2014). The use of eDNA to detect biodiversity shifts has been an active area of research (Lodge *et al.*, 2012; Kelly *et al.*, 2014; Kelly, 2016). Metabarcoding is also emerging as an invaluable tool in the examination of trophic relationships, through HTS analysis of gut contents and fecal material (Deagle *et al.*, 2013; Albaina *et al.*, 2016). Both these applications require specific examination of the impact of degraded DNA on the accuracy and reliability of the analyses, since amplicon size and copy number will impact metabarcoding uncertainty biases and will need to be addressed prior to use in ecological monitoring.

Metabarcoding may also substantially improve capabilities for accurate identification and early detection of introduced NIS (Mountfort *et al.*, 2012; Kelly *et al.*, 2014;

Zaiko *et al.*, 2015b). Such early-warning will provide managers with options to act before a harmful species can achieve high abundance (Robinson *et al.*, 2011). Shipping, in particular transported ballast water, is considered to be one of the most important pathways of marine biological invasions worldwide (Molnar *et al.*, 2008), yet traditional sampling does not always capture all organisms, especially at the early phase of invasion (Lehtiniemi *et al.*, 2015). Previous studies have shown that estimates of taxon-specific DNA concentrations determined using qPCR correlate positively with abundance estimates of that taxon (Thomsen *et al.*, 2012). Clearly, accurate detection and quantification are important for managers to determine the phase of the invasion and necessary approaches for eradication.

A promising use of metabarcoding for management of NIS concerns Ballast Water Management Convention compliance control. Metabarcoding can be used to verify positive results in control surveys (i.e. zero counts of organisms by microscopic analyses), because some species can be overlooked by conventional analysis. Metabarcoding, especially when used in combination with morphological analyses (Zaiko *et al.*, 2015a), is a powerful new tool for NIS monitoring and management.

SUMMARY AND RECOMMENDATIONS

Metabarcoding is revolutionizing the analysis of marine biodiversity and has a significant advantage of detecting the 'hidden diversity' of marine zooplankton (Lindeque *et al.*, 2013). Emerging results indicate that estimates of global zooplankton diversity will markedly increase with more accurate definition and higher resolution of time/space patterns made possible by metabarcoding. In addition, metabarcoding will allow more rapid detection and description of the impacts of climate change on biodiversity and biogeography. As HTS becomes more accessible and less expensive, the use of metabarcoding will expand into numerous applications in ocean monitoring and management, including calculation of biotic indices, trophic interactions and food web analysis and detection of introduced NIS. Among the challenges remaining for reliable and routine application of metabarcoding for analysis of zooplankton are evaluation and comparison of results using various barcode gene regions (as well as different primers and protocols); development of methodologies using more variable gene regions that can ensure identification, discrimination and detection of closely related, cryptic and rare species; impacts of degraded DNA (e.g. environmental DNA and DNA recovered from gut contents) and continued development of taxonomic-ally comprehensive reference databases for all gene

regions. A particular need is to move metabarcoding applications from identification and detection of taxa to their quantification in terms of abundance and/or biomass, which will require concerted effort to address biases associated with gene CNV. Despite the remarkable promise of metabarcoding in yielding new understanding and appreciation for global patterns of zooplankton diversity, it is critically important to maintain expertise and capacity in morphological taxonomic identification of zooplankton to ensure that metabarcoding approaches can be validated. Such integrative morphological and molecular taxonomic approaches will provide the necessary foundation and future of research, monitoring and management of the pelagic realm.

ACKNOWLEDGMENTS

This manuscript results from discussions of the International Council for the Exploration of the Sea (ICES) Working Group on Integrated Morphological and Molecular Taxonomy (WGIMT; see <http://www.ices.dk/community/groups/Pages/WGIMT.aspx> and <http://wgimt.net/>). The authors gratefully acknowledge all members of WGIMT for their contribution to the development of the ideas presented here. This paper benefitted from collaboration with ICES WGIMTMO and WGBOSV. This is contribution number 748 of the Marine Research Division of AZTI.

FUNDING

We acknowledge support from the International Council for the Exploration of the Sea (ICES) for WGIMT, as well as conferences and symposia related to the Working Group's mission. ICES Delegates from member countries provided financial support to some WGIMT members for travel to annual meetings. Additional support was provided by: NOAA Office of Ocean Exploration and Research (Grant No. NA15OAR0110209 to AB, University of Connecticut, USA); UK Natural Environment Research Council and Department for Environment, Food and Rural Affairs (Grant No. NE/L003279/1 to PKL, Plymouth Marine Laboratory, UK); DEVOTES (DEVELOPMENT OF innovative TOOLS for understanding marine biodiversity and assessing Good Environmental Status) project funded by the European Union (EU) 7th Framework Program (Grant No. 308392 to NR-E, AZTI, Spain); Basque Government (Grant No. IT558-10 to AA, University of the Basque Country, UPV/EHU, Spain); and BONUS BIO-C3 project and BONUS (Art 185), funded by the EU and Finnish Academy (Grant No. 273660 to ML, Finnish Environment Institute, Finland).

REFERENCES

- Albaina, A., Aguirre, M., Abad, D., Santos, M. and Estonba, A. (2016) 18S rRNA V9 metabarcoding for diet characterization: a critical evaluation with two sympatric zooplanktivorous fish species. *Ecol. Evol.*, doi: 10.1002/ece3.1986.
- Amaral-Zettler, L. A., McCliment, E. A., Ducklow, H. W. and Huss, S. M. (2009) A method for studying protistan diversity using massively parallel sequencing of V9 hypervariable regions of small-subunit ribosomal RNA genes. *PLoS One*, **4**, e6372.
- Amend, A. S., Seifert, K. A. and Bruns, T. D. (2010) Quantifying microbial communities with 454 pyrosequencing: does read abundance count? *Mol. Ecol.*, **19**, 5555–5565.
- Aylagas, E., Borja, A. and Rodríguez-Ezpeleta, N. (2014) Environmental status assessment using DNA metabarcoding: towards a genetics based marine biotic index (gAMBI). *PLoS One*, **9**, e90529.
- Beaugrand, G., Edwards, M. and Legendre, L. (2010) Marine biodiversity, ecosystem functioning, and carbon cycles. *Proc. Natl Acad. Sci. USA*, **107**, 10120–10127.
- Bik, H. M., Porazinska, D. L., Creer, S., Caporaso, J. G., Knight, R. and Thomas, W. K. (2012) Sequencing our way towards understanding global eukaryotic biodiversity. *Trends Ecol. Evol.*, **27**, 233–243.
- Bohmann, K., Evans, A., Gilbert, M. T., Carvalho, G. R., Creer, S., Knapp, M., Yu, D. W. and de Bruyn, M. (2014) Environmental DNA for wildlife biology and biodiversity monitoring. *Trends Ecol. Evol.*, **29**, 358–367.
- Bourlat, S. J., Borja, A., Gilbert, J., Taylor, M. I., Davies, N., Weisberg, S. B., Griffith, J. F., Lettieri, T. *et al.* (2013) Genomics in marine monitoring: new opportunities for assessing marine health status. *Mar. Pollut. Bull.*, **74**, 19–31.
- Brown, E. A., Chain, F. J. J., Crease, T. J., MacIsaac, H. J. and Cristescu, M. E. (2015) Divergence thresholds and divergent biodiversity estimates: can metabarcoding reliably describe zooplankton communities? *Ecol. Evol.*, **5**, 2234–2251.
- Bucklin, A., Hopcroft, R. R., Kosobokova, K. N., Nigro, L. M., Ortman, B. D., Jennings, R. M. and Wiebe, P. H. (2010a) DNA barcoding of Arctic Ocean holozooplankton for species identification and recognition. *Deep Sea Res. II*, **57**, 40–48.
- Bucklin, A., Ortman, B. D., Jennings, R. M., Nigro, L. M., Sweetman, C. J., Copley, N. J. and Sweetman, C. J. (2010b) A “Rosetta Stone” for metazoan zooplankton: DNA barcode analysis of species diversity of the Sargasso Sea (Northwest Atlantic Ocean). *Deep Sea Res. II*, **57**, 2234–2247.
- Bucklin, A., Steinke, D. and Blanco-Bercial, L. (2011) DNA barcoding of marine metazoa. *Annu. Rev. Mar. Sci.*, **3**, 471–508.
- Carugati, L., Corinaldesi, C., Dell’Anno, A. and Danovaro, R. (2015) Metagenetic tools for the census of marine meiofaunal biodiversity: an overview. *Mar. Genomics*, doi:10.1016/j.margen.2015.04.010.
- Corell, J. and Rodríguez-Ezpeleta, N. (2014) Tuning of protocols and marker selection to evaluate the diversity of zooplankton using metabarcoding. *Rev. Invest. Mar.*, **21**, 19–39.
- Deagle, B. E., Thomas, A. C., Shaffer, A. K., Trites, A. W. and Jarman, S. N. (2013) Quantifying sequence proportions in a DNA-based diet study using Ion Torrent amplicon sequencing: which counts count? *Mol. Ecol. Res.*, **13**, 620–633.
- DeVargas, C., Audic, S., Henry, N., Decelle, J., Mahé, F., Logares, R., Lara, E., Berney, C. *et al.* (2015) Eukaryotic plankton diversity in the sunlit ocean. *Science*, **348**, 1261605.

- Dowle, E. J., Pochon, X., Banks, J. C., Shearer, K. and Wood, S. A. (2015) Targeted gene enrichment and high-throughput sequencing for environmental biomonitoring: a case study using freshwater macroinvertebrates. *Mol. Ecol. Res.*, doi: 10.1111/1755-0998.12488.
- Eiler, A., Drakare, S., Bertilsson, S., Pernthaler, J., Peura, S., Rofner, C., Simek, K., Yang, Y. *et al.* (2013) Unveiling distribution patterns of freshwater phytoplankton by a next generation sequencing based approach. *PLoS One*, **8**, e53516.
- Floyd, R., Abebe, E., Papert, A. and Blaxter, M. (2002) Molecular barcodes for soil nematode identification. *Mol. Ecol.*, **11**, 839–850.
- Fonseca, V. G., Carvalho, G. R., Sung, W., Johnson, H. F., Power, D. M., Neill, S. P., Packer, M., Blaxter, M. L. *et al.* (2010) Second-generation environmental sequencing unmasks marine metazoan biodiversity. *Nat. Commun.*, **1**, 98.
- Goetze, E. (2010) Species discovery in marine planktonic invertebrates through global molecular screening. *Mol. Ecol.*, **19**, 952–967.
- Hebert, P. D. N., Ratasingham, S. and DeWaard, J. R. (2003) Barcoding animal life: cytochrome c oxidase subunit 1 divergences among closely related species. *Proc. R. Soc. Lond. B Biol.*, **270**, S96–S99.
- Hirai, J., Kuriyama, M., Ichikawa, T., Hidaka, K. and Tsuda, A. (2015a) A metagenetic approach for revealing community structure of marine planktonic copepods. *Mol. Ecol. Res.*, **15**, 68–80.
- Hirai, J., Shimode, S. and Tsuda, A. (2013) Evaluation of ITS2-28S as a molecular marker for identification of calanoid copepods in the subtropical western North Pacific. *J. Plankton Res.*, **35**, 644–656.
- Hirai, J. and Tsuda, A. (2015) Metagenetic community analysis of epipelagic planktonic copepods in the tropical and subtropical Pacific. *Mar. Ecol. Progr. Ser.*, **534**, 65–78.
- Hirai, J., Yasuike, M., Fujiwara, A., Nakamura, Y., Hamaoka, S., Katakura, S., Takano, Y. and Nagai, S. (2015b) Effects of plankton net characteristics on metagenetic community analysis of metazoan zooplankton in a coastal marine ecosystem. *J. Exptl. Mar. Biol. Ecol.*, **469**, 36–43.
- Ji, Y., Ashton, L., Pedley, S. M., Edwards, D. P., Tang, Y., Nakamura, A., Kitching, R., Dolman, P. M. *et al.* (2013) Reliable, verifiable and efficient monitoring of biodiversity via metabarcoding. *Ecol. Lett.*, **16**, 1245–1257.
- Kelly, R. P. (2016) Making environmental DNA count. *Mol. Ecol. Res.*, **16**, 10–12.
- Kelly, R. P., Port, J. A., Yamahara, K. M., Martone, R. G., Lowell, N., Thomsen, P. F., Mach, M. E., Prahler, E. *et al.* (2014) Harnessing DNA to improve environmental management. *Science*, **344**, 1455–1456.
- Lehtiniemi, M., Ojaveer, H., David, M., Galil, B., Gollasch, S., McKenzie, C., Minchin, D., Occhipinti-Ambrogi, A. *et al.* (2015) Dose of truth—monitoring marine non-indigenous species to serve legislative requirements. *Mar. Policy*, **54**, 26–35.
- Lindeque, P. K., Harris, R. P., Jones, M. B. and Smerdon, G. R. (1999) Simple molecular method to distinguish the identity of *Calanus* species (Copepoda: Calanoida) at any developmental stage. *Mar. Biol.*, **133**, 91–96.
- Lindeque, P. K., Hay, S. J., Heath, M. R., Invarsdottir, A., Rasmussen, J., Smerdon, G. R. and Waniek, J. J. (2006) Merging conventional microscopy and molecular analysis to analyse the species composition of plankton samples. *J. Plankton Res.*, **28**, 221–238.
- Lindeque, P. K., Parry, H. E., Harmer, R. A., Somerfield, P. J. and Atkinson, A. (2013) Next generation sequencing reveals the hidden diversity of zooplankton assemblages. *PLoS One*, **8**, e81327.
- Lindsay, D. J., Grossman, M. M., Nishikawa, J., Bentlage, B. and Collins, A. G. (2015) DNA barcoding of pelagic cnidarians: current status and future prospects. *J. Plankton Soc. Japan*, **62**, 39–43.
- Lodge, D. M., Turner, C. R., Jerde, C. L., Barnes, M. A., Chadderton, L., Egan, S. P., Feder, J. L., Mahon, A. R. *et al.* (2012) Conservation in a cup of water: estimating biodiversity and population abundance from environmental DNA. *Mol. Ecol.*, **21**, 2555–2558.
- Machida, R., Hashiguchi, Y., Nishida, M. and Nishida, S. (2009) Zooplankton diversity analysis through single-gene sequencing of a community sample. *BMC Genomics*, **10**, 438.
- Mallatt, J. M., Garey, J. R. and Shultz, J. W. (2004) Ecdysozoan phylogeny and Bayesian inference: first use of nearly complete 28S and 18S rRNA gene sequences to classify the arthropods and their kin. *Mol. Phylogenet. Evol.*, **31**, 178–191.
- Massana, R., Gobet, A., Audic, S., Bass, D., Bittner, L., Boutte, C., Chambouvet, A., Christen, R. *et al.* (2015) Marine protist diversity in European coastal waters and sediments as revealed by high-throughput sequencing. *Environ. Microbiol.*, doi:10.1111/1462-2920.12955.
- Mohrbeck, I., Raupach, M. J., Martínez Arbizu, P., Knebelberger, T. and Laakmann, S. (2015) High throughput sequencing—the key to rapid biodiversity assessment of marine metazoa? *PLOS One*, **10**, e0140342.
- Möllmann, C. and Diekmann, R. (2012) Marine ecosystem regime shifts induced by climate and overfishing: a review for the Northern Hemisphere. *Adv. Ecol. Res.*, **47**, 303.
- Molnar, J. L., Gamboa, R. L., Revenga, C. and Spalding, M. D. (2008) Assessing the global threat of invasive species to marine biodiversity. *Front. Ecol. Environ.*, **6**, 485–492.
- Mountfort, D., Smith, K. F., Kirs, M., Kuhajek, J., Adamson, J. E. and Wood, S. A. (2012) Development of single and multispecies detection methods for the surveillance and monitoring of marine pests in New Zealand. *Aquat. Invas.*, **7**, 125–128.
- Pearman, J. K., El-Sherbiny, M. M., Lanzén, A., Al-Aidaros, A. M. and Irigoien, X. (2014) Zooplankton diversity across three Red Sea reefs using pyrosequencing. *Front. Mar. Sci.*, **1**, doi:10.3389/fmars.2014.00027.
- Pearman, J. K. and Irigoien, X. (2015) Assessment of zooplankton community composition along a depth profile in the central Red Sea. *PLoS ONE*, **10**, e0133487.
- Prokopowich, C. D., Gregory, T. R. and Crease, T. J. (2003) The correlation between rDNA copy number and genome size in eukaryotes. *Genome*, **46**, 48–50.
- Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J. and Glöckner, F. O. (2013) The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucl. Acids Res.*, **41**, D590–D596.
- Robinson, A., Burgman, M. A. and Cannon, R. (2011) Allocating surveillance resources to reduce ecological invasions: maximizing detections and information about the threat. *Ecol. Appl.*, **21**, 1410–1417.
- Shokralla, A., Spall, J. L., Gibson, J. F. and Hajibabaei, M. (2012) Next-generation sequencing technologies for environmental DNA research. *Mol. Ecol.*, **21**, 1794–1805.
- Sun, C., Zhao, Y., Li, H., Dong, Y., MacIsaac, H. J. and Zhan, A. (2015) Unreliable quantitation of species abundance based on high-

- throughput sequencing data of zooplankton communities. *Aquat. Biol.*, **24**, 9–15.
- Tang, C. Q., Leasi, F., Obertegger, U., Kieneke, A., Barraclough, T. G. and Fontaneto, D. (2012) The widely used small subunit 18S rDNA molecule greatly underestimates true diversity in biodiversity surveys of the meiofauna. *Proc. Natl Acad. Sci. USA*, **109**, 16208–16212.
- Thomsen, P. F., Kielgast, J., Iversen, L. L., Wiuf, C., Rasmussen, M., Gilbert, M. T., Orlando, L. and Willerslev, E. (2012) Monitoring endangered freshwater biodiversity using environmental DNA. *Mol. Ecol.*, **11**, 2565–2573.
- Zaiko, A., Martínez, J. L., Schmidt-Petersen, J., Ribicic, D., Samuiloviene, A. and Garcia-Vazquez, E. (2015a) Metabarcoding approach for the ballast water surveillance—an advantageous solution or an awkward challenge? *Mar. Pollut. Bull.*, **92**, 25–34.
- Zaiko, A., Samuiloviene, A., Ardura, A. and Garcia-Vazquez, E. (2015b) Metabarcoding approach for nonindigenous species surveillance in marine coastal waters. *Mar. Pollut. Bull.*, **100**, 53–59.
- Zhan, A. and MacIsaac, H. J. (2015) Rare biosphere exploration using high-throughput sequencing: research progress and perspectives. *Conserv. Genet.*, **16**, 513–522.
- Zheng, L., He, J., Lin, Y., Cao, W. and Zhang, W. (2014) 16S rRNA is a better choice than COI for DNA barcoding hydrozoans in the coastal waters of China. *Acta Oceanol. Sin.*, **33**, 55–76.